

Alphabetical Statistical Symbols:

| Symbol | Text Equivalent | Meaning | Formula | Link to Glossary (if appropriate) |
|------------|---|---|---|-----------------------------------|
| a | | Y- intercept of least square regression line | $a = \bar{y} - b\bar{x}$, for line $y = a + bx$ | Regression: y on x |
| b | | Slope of least squares regression line | $b = \frac{\sum (x - \bar{x})(y - \bar{y})}{\sum (x - \bar{x})^2}$ for line $y = a + bx$ | Regression: y on x |
| B (n, p) | Binomial distribution with parameters n and p | Discrete probability distribution for the probability of number of successes in n independent random trials under the identical conditions. | If X follows B (n, p) then, $P(X = r) = {}^n C_r p^r (1 - p)^{n-r}$, Where, $0 < p < 1$, $r = 0, 1, 2, \dots, n$ | Binomial Distribution |
| c | | Confidence level | $c = P(-z_c < Normal(0,1) < z_c)$ | Confidence interval |
| ${}^n C_r$ | n-c-r | Combinations (number of combinations of n objects taken r at a time) | ${}^n C_r = \frac{n!}{r!(n-r)!}$, where $n \geq r$ | |
| $C_{n,r}$ | n-c-r | Combinations (number of combinations of n objects taken r at a time) | $C_{n,r} = \frac{n!}{r!(n-r)!}$, where $n \geq r$ | |
| Cov (X, Y) | Covariance between X and Y | Covariance between X & Y | $Cov(X) = E[(X - E(X))(Y - E(Y))]$ | |

100+ online courses in statistics

| Symbol | Text Equivalent | Meaning | Formula | Link to Glossary (if appropriate) |
|----------------|-------------------------|---------------------------|---|---|
| CV | | Coefficient of variation | $CV = \frac{\text{Standard Deviation}}{\text{Arithmetic mean}}$ | |
| df | | Degree(s) of freedom | | |
| E | | Maximal error tolerance | $E = z_c \frac{\sigma}{\sqrt{n}}$ for large samples. | |
| E (f (x)) | Expected value of f (x) | | $E (f (x)) = \sum f(x)P(x)$ | |
| f | | Frequency | f = number of times score. | |
| F | | F-distribution variable | $F = \frac{\chi_1^2 / n_1}{\chi_2^2 / n_2}$ where n_1 and n_2 are the corresponding degrees of freedom. | F-distribution, Hypothesis testing for equality of 2 variances. |
| F (x) or F_x | | Distribution function | $F_x = \int_{-\infty}^x f_x dx$ | |
| f (x) or f_x | | Probability mass function | Depends on the distribution. $f_x \geq 0$ & $\int_x f_x dx = 1$. | |
| H_0 | H-naught | Null hypothesis. | The null hypothesis is the hypothesis about the population parameter. | Testing of hypothesis |
| H_1 | H-one | Alternate hypothesis. | An alternate hypothesis is constructed in such a way that it is the one to be accepted when the null hypothesis must be rejected. | Testing of hypothesis |
| IQR | | Interquartile range | $IQR = Q_3 - Q_1$ | Measures of central tendency. |

100+ online courses in statistics

| Symbol | Text Equivalent | Meaning | Formula | Link to Glossary (if appropriate) |
|------------|--------------------------|--|---|-----------------------------------|
| MS | M-S | Mean square | $MS = \frac{SS}{df}$ | Analysis of variance (ANOVA) |
| n | | Sample size. | n = number of units in a sample. | |
| N | | Population size | N = Number of units in the population. | |
| $P_{n,r}$ | n-p-r | Permutation (number of ways to arrange in order n distinct objects taking them r at a time) | $P_{n,r} = \frac{n!}{(n-r)!}$, where $n \geq r$ | |
| ${}_n P_r$ | n-p-r | Permutation (number of ways to arrange in order n distinct objects taking them r at a time) | ${}_n P_r = \frac{n!}{(n-r)!}$, where $n \geq r$ | |
| \hat{p} | p-hat | Sample proportion | $\hat{p} = \frac{\text{number of success}}{\text{number of trials}}$ | Binomial distribution |
| P (A B) | Probability of A given B | Conditional probability | $P(A B) = \frac{P(A \cap B)}{P(B)}$ | |
| P (x) | Probability of x | Probability of x | $P(x) = \frac{\text{No.of favorable outcomes}}{\text{Total no.of outcomes}}$ | |
| p-value | | The attained level of significance. | P value is the smallest level of significance for which the observed sample statistic tells us to reject the null hypothesis. | |
| Q | | Probability of not happening of the event | $q = 1 - p$ | |

| Symbol | Text Equivalent | Meaning | Formula | Link to Glossary (if appropriate) |
|---------|-----------------|----------------------------------|--|-----------------------------------|
| Q_1 | Q-one | First quartile | Q_1 = Median of the lower half of the data that is data below median. | Measures of central tendency |
| Q_2 | Q-two | Second quartile Or Median | Q_2 = Central value of an ordered data. | Measures of central tendency |
| Q_3 | Q-three | Third quartile | Q_3 = Median of the upper half of the data that is data above the median. | Measures of central tendency |
| R | | Sample Correlation coefficient | $r = \frac{Co\ variance(X,Y)}{[SD(X)]*[SD(Y)]}$ | |
| r^2 | r-square | Coefficient of determination | $r^2 = (Correlation\ coefficient)^2$ | |
| R^2 | r-square | Multiple correlation coefficient | $R^2 = 1 - \frac{mean\ square\ error}{S_y^2}$ | |
| S | | Sample standard deviation | $s = \sqrt{\frac{\sum (x - \bar{x})^2}{n - 1}}$ for ungrouped data. $s = \sqrt{\frac{\sum f(x - \bar{x})^2}{(\sum f) - 1}}$ for grouped data. | Measures of dispersion |
| S^2 | S-square | Sample variance | $S^2 = \frac{\sum (x - \bar{x})^2}{n - 1}$ for ungrouped data. $S^2 = \frac{\sum f(x - \bar{x})^2}{(\sum f) - 1}$ for grouped data | Measures of dispersion |
| S_e^2 | s-e- square | Error variance | $S_e^2 = \frac{sum\ of\ squares\ of\ residuals}{n}$ | |

100+ online courses in statistics

| Symbol | Text Equivalent | Meaning | Formula | Link to Glossary (if appropriate) |
|-----------------|-----------------|---|--|-----------------------------------|
| SD | | Sample standard deviation | $s = \sqrt{\frac{\sum (x - \bar{x})^2}{n - 1}}$ for ungrouped data. $s = \sqrt{\frac{\sum f(x - \bar{x})^2}{(\sum f) - 1}}$ for grouped data. | |
| sk _b | | Bowley's coefficient of skewness | $sk_b = \frac{(Q_3 - Q_2) - (Q_2 - Q_1)}{(Q_3 - Q_1)}$ | Measures of skewness |
| sk _p | | Pearson's coefficient of skewness | $sk_p = \frac{\text{Mean} - \text{Mode}}{\text{Standard Deviation}}$ | Measures of skewness |
| SS _x | | Sum of Squares | $SS_x = \sum (x - \bar{x})^2$ for ungrouped data. $SS_x = \sum f(x - \bar{x})^2$ for grouped data. | |
| t | | Student's t variable. | $t = \frac{\text{Normal}(0,1)}{\sqrt{\chi_n^2/n}}$ | t-distribution |
| t _c | t critical | The critical value for a confidence level c. | t _c = Number such that the area under the t distribution for a given number of degrees of freedom falling between -t _c and t _c is equal to c. | Testing of hypothesis |
| Var (X) | Variance of X | Variance of X | Var (X) = E (X - μ) ² | |
| X | | Independent variable or explanatory variable in regression analysis | Eg. In the study of, yield obtained & the irrigation level, independent variable is, X= Irrigation level. | |

100+ online courses in statistics

| Symbol | Text Equivalent | Meaning | Formula | Link to Glossary (if appropriate) |
|-----------|-----------------|---|---|--|
| \bar{x} | x-bar | Arithmetic mean or Average of X scores. | $\bar{x} = \frac{\sum x}{n}$ for ungrouped data. $\bar{x} = \frac{\sum fx}{\sum f}$ for grouped data. | Measures of central tendency |
| y | | Dependent variable or response variable in regression analysis | Eg. In the study of, yield obtained & the irrigation level, dependent variable is, Y= Yield obtained. | |
| Z | Z-score | Standard normal variable (Normal variable with mean = 0 & SD = 1) | $z = \frac{x - \mu}{\sigma}$, where X follows Normal (μ, σ). | Standard normal distribution |
| z_c | z critical | The critical value for a confidence level c. | z_c = Number such that the area under the standard normal curve falling between $-z_c$ and z_c is equal to c. | Testing of hypothesis Confidence interval |

Greek Statistical Symbols:

| Symbol | Text Equivalent | Meaning | Formula | Link to Glossary (if appropriate) |
|----------|-----------------|--|---|-----------------------------------|
| α | Alpha | Type I error or Level of Significance. | $\alpha = P$ [Rejecting the null hypothesis Null hypothesis is true]. | Hypothesis Testing |

100+ online courses in statistics

| Symbol | Text Equivalent | Meaning | Formula | Link to Glossary (if appropriate) |
|-------------|-----------------|---|--|-----------------------------------|
| β | Beta | Type II error or Power of the test. | $\beta = P$ [Accepting the null hypothesis Null hypothesis is False]. | Hypothesis Testing |
| ϵ | Epsilon | “Error Term” in regression/statistics; more generally used to denote an arbitrarily small positive number | $y = \beta_0 + \beta_1 *x + \epsilon$ | Regression |
| χ^2 | Chi-square | Chi-square distribution | $\chi^2 =$ Sum of n independent Standard normal variables | Chi-square distribution. |
| χ^2 | Chi-square | Chi-square distribution | $\chi^2 = \sum \frac{(O - E)^2}{E}$ where O is the observed frequency and E is the expected frequency. Or $\chi^2 = \frac{(n-1)s^2}{\sigma^2}$ (?) | Goodness of fit test |
| $\Gamma(n)$ | Gamma-n | Gamma function | $\Gamma(n) = (n-1) !$ | |
| λ | Lambda | Parameter used for Poisson distribution | $\lambda =$ Mean of Poisson distribution | Poisson distribution |
| μ | Mu | Arithmetic mean or Average of the population. | $\mu = \frac{\sum x}{N}$ $\mu = E(x) = \sum xP(x)$ | |
| μ_r | Mu-r | r^{th} central moment | $\mu_r = E [(X - \mu)^r]$ | Measures of central tendency. |
| μ_r' | Mu-r-dash | r^{th} Raw moment | $\mu_r' = E (X^r)$ | Measures of central tendency. |
| ρ | Rho | Population correlation coefficient | $\rho = \frac{\text{Covariance}(X, Y)}{SD(X) * SD(Y)}$ | |

100+ online courses in statistics

| Symbol | Text Equivalent | Meaning | Formula | Link to Glossary (if appropriate) |
|------------|-----------------|-------------------------------|--|-----------------------------------|
| Σ | Sigma | Summation | $\Sigma x = \text{Sum of } x \text{ scores.}$ | |
| σ | Sigma | Population Standard Deviation | $\sigma = \sqrt{\frac{\Sigma (x - \mu)^2}{N}}$ $\sigma = \sqrt{E[(x - \mu)^2]} = \sqrt{\Sigma (x - \mu)^2 P(x)}$ | Measures of dispersion |
| σ^2 | Sigma square | Population variance | $\sigma^2 = \frac{\Sigma (x - \mu)^2}{N}$ | Measures of dispersion |

Mathematical Statistical Symbols:

| Symbol | Text Equivalent | Meaning | Formula | Link to Glossary (if appropriate) |
|--------|-----------------|--|---|-----------------------------------|
| ! | Factorial | Product of all integers up to the given number | $n! = n (n-1) (n-2) \dots\dots\dots 1.$ $0! = 1$ | |
| c | Complement | not | For example: A^c is not A | |
| \cup | Union | or | For example: $(A \cup B)$ is happening of either event A or event B | |
| \cap | Intersection | And | For example: $(A \cap B)$ is happening of both event A and event B | |

100+ online courses in statistics